

3-200 keV spectral states and variability of the INTEGRAL Black Hole binary IGR J17464-3213¹

F. Capitanio², P. Ubertini², A. Bazzano², P. Kretschmar^{3,4}, A. A. Zdziarski⁵, A. Joinet⁶,
E.J. Barlow⁷, A.J.Bird⁷, A.J.Dean⁷, E. Jourdain⁶, G. De Cesare², M. Del Santo², L.
Natalucci², M.Cadolle Bel⁸, A. Goldwurm⁸

ABSTRACT

On March 2003, IBIS, the gamma-ray imager on board the *INTEGRAL* satellite, detected an outburst from a new source, IGR J17464-3213, that turned out to be a *HEAO-1* transient, H1743-322. In this paper we report on the high energy behaviour of this BHC studied with the three main instruments onboard *INTEGRAL*. The data, collected with unprecedented sensitivity in the hard X-Ray range, show a quite hard Comptonised emission from 3 keV up to 150 keV during the rising part of the source outburst, with no thermal emission detectable. A few days later, a prominent soft disk multicolour component appears, with the hard tail luminosity almost unchanged: $\sim 5 \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$. Two months later, during a second monitoring campaign near the end of the outburst, the observed disk component was unchanged. Conversely, the Comptonised emission from the central-hot part of the disk reduced by a factor of ~ 10 . We present here its long term behaviour in different energy ranges and the combined JEM-X, SPI and IBIS wide band spectral evolution of this source.

Subject headings: Galaxy:High Energy — gamma rays: LMXB, Black Hole Candidates IGR J17464-3213

¹Based on observations with *INTEGRAL*, an ESA project with instruments and science data centre funded by ESA member states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), Czech Republic and Poland and with the participation of Russia and the USA.

²IASF, CNR-INAF, Rome, Italy, Via Fosso del Cavaliere 100, I-00133 Rome, Italy

³MPE, Postfach 1603, 85740 Garching, Germany

⁴*INTEGRAL* Science Data Centre, Chemin d'Ecogia 16, 1291, Versoix, Switzerland

⁵Centrum Astronomiczne im. M. Kopernika, 18 00716 Warszawa, Poland

⁶CESR, Toulouse, France

⁷School of Physics and Astronomy, University of Southampton, Highfield, Southampton, SO 17 1BJ, UK.

⁸Sap-CEA, Saclay, F-91191 Gif-sur-Yvette, France

1. Introduction

On 2003 March 21 during the Galactic Centre Deep Exposure (GCDE) Program *INTEGRAL* detected a relatively bright source (~ 60 mCrab at 15-40 keV) named IGR J17464-3213 (Revnivtsev et al. 2003). The source was then localised at R.A. (2000) = $17^{\text{h}}46.3^{\text{m}}$, Dec. = $-32^{\circ}14.4'$, with an error box of 1.6 arcmin (90% confidence) and associated with H1743-322 (Markwardt et al. 2003), a bright BHC observed by HEAO1 in 1977 with an intensity of 700 mCrab at 2-10 keV and localised with two possible positions (Doxsey et al. 1977). The error box of IGR J17464-3213 was compatible with only one of these two, thereby resolving this 25 year old ambiguity. After the 2003 outburst, follow up observations with *RXTE* and *INTEGRAL* reported strong flux and possible spectral variability (Parmar et al. 2003). *RXTE* observed the source for the first time on 2003 March 29 during a PCA Galactic bulge scan followed by a pointed observation. The mean PCA fluxes, in the bands 2-10, 15-40 and 40-100 keV, were 50, 200, 220 mCrab respectively. The spectrum was consistent with an absorbed power law with a photon index 1.49 ± 0.02 , and a column density: $N_{\text{H}} = 2.4 \times 10^{22} \text{ cm}^{-2}$ (Markwardt et al. 2003). In this paper we focus on the data collected with the coded mask co-aligned telescopes IBIS (15 keV - 10 MeV) (Ubertini et al. 2003), SPI (20 keV - 8 MeV) (Vedrenne et al. 2003) and JEM-X (3-35 keV) (Lund et al. 2003) on board *INTEGRAL* (Winkler et al. 2001). The source was monitored by *INTEGRAL* in the framework of the Core Programme (CP) observations (Kretschmar et al. 2003; Grebenev et al. 2003). The data analysed here cover the period from the beginning of the outburst (March 2003) to its end (October 2003).

2. Data Analysis

The analysed data set consists of all CP observations in which IGR J17464-3213 was within the high energy detectors field of view. In particular, the observations were performed in three different periods: from 2003 March 12 to 2003 April 15, from 2003 August 19 to 20, from 2003 October 4 to 9. The first period contains four observations of the pre-outburst state of the source; the other two shorter observations cover the end of the outburst phase. All the CP observations are organised into un-interrupted 2000s long science windows (SCW): light curves, hardness ratio and spectra are then extracted for each individual SCW. Wide band spectra of the source are obtained using data from the three high energy instruments (JEM-X, IBIS and SPI). In particular, the IBIS data were processed using the Off-line Scientific Analysis (OSA version 3) (Goldwurm et al. 2003) released by the *INTEGRAL* Scientific Data Centre (ISDC) (Courvoisier et al. 2003) with a modified pipeline in order to optimise the source spectral extraction. The IBIS data set has been divided in two subsets

according to the source position being within the partially coded field of view (PCFOV) or in the narrower fully coded field of view (FCFOV). Both data sets have been used, after source intensity correction for off-axis detectors response, to produce the source light curves in different energy ranges. The off-axis correction was made using the available calibration data, the accuracy on the flux being from 10% at low energy (15-20 keV) up to 5% at higher energy ($> 60\text{keV}$). Only FCFOV data, less affected by the off-axis response of the gamma-ray instruments, was selected for spectral extraction and fitting.

SPI spectra were extracted using software specially developed for fitting the positions and the fluxes of all significant sources in the field of view. The background model used for SPI data is based on an uniformity map determined for each energy band from empty fields observation.

JEM-X spectra were derived with the ISDC OSA version 4 release, and updated spectral matrices. While IBIS and SPI provide a very large FOV ($>30\text{ deg}$), Jem-X has a narrower FOV ($>10\text{ deg}$), thus providing only a partial overlap with the high energy detectors. When spectral data are obtained in the same time interval from more then one instrument, the proper normalizing constant is added in the data fit, taking into account the best knowledge of the instrument cross calibration. In optimum conditions, the *INTEGRAL* spectra we produced cover the range from 3 keV to 200 keV.

3. Results

3.1. The time evolution of the outburst

The source was continuously monitored by the *RXTE* ASM and whenever possible by *INTEGRAL* as shown in Figure 1, with the simultaneous *INTEGRAL* observation periods represented by dashed rectangles. The monitoring of the source was quite continuous during the pre-outburst episode (from revolution 53 to 61) and then only two deep observations were performed after about 4 (revolution 103) and 5.5 month (revolutions 119-120). In our work we have not analysed the data of the outburst itself as they are proprietary data published elsewhere (Parmar et al. 2003). We have produced light curves in the 20-30, 30-40, 40-80 keV energy ranges and investigated the Hardness Ratio, defined as $HR = (flux_{40-80} - flux_{20-30}) / (flux_{40-80} + flux_{20-30})$. The IBIS 20-30 keV and 40-60 keV light curves of the initial part of the outburst, flux averaged over one SCW, are shown in Figure 2. The IBIS light curves show strong variations and it is possible to note several short peaks in the flux intensity with different shapes in the lower and higher energy bands. From the beginning of the IBIS observation, MJD ~ 52729 to ~ 52731 , (Modified Julian Date) the source averaged intensity increases monotonically in the lower energy range while the flux is

almost unchanged at higher energies: this is reflected in the anticorrelation shown by the HR that decreases continuously in the same week as can be seen in Figure 2(c). The few points sampled in the next 20 days, due to a limited coverage, show a decrease in both low energy flux and HR. Starting from day ~ 52744 , the source shows a clearly visible flux increase in a few hours by a factor of 2.6, 3.0 and 3.2 in the 20-30, 30-40 and 40-60 keV ranges respectively, without any corresponding hardening. After the peak, the source emission falls by a factor of ~ 4.5 in the following hours (20-30 keV) with a small decrease of the HR; the soft part of the spectrum becomes now more prominent (Figure 2). This corresponds to a dip in the flux emission, lasting a few hours (around day 52745.1), during which the source shows the lowest observed value for the HR. Finally, the high energy component, and the HR, increase again (MJD=52745.5). The source shows a similar behaviour around MJD=52750. To look for a possible correlation of the Flux vs. Spectral index we have fitted the available IBIS spectra during revolution 53, 57, 59-61 and 119-120, with a Cutoff power law model. We noticed that the cut off energy of the model had substantially the same value in each fit, so we freeze it to an average value of $E=53.3$ keV. The reduced spectral fit χ^2 covers a range from 0.8 to 1.3. The result, summarised in Figure 3, shows that the spectral index has a “circular” or “hysteresis” like behaviour. In fact, at the end of the main outburst (revolution 103-120), the source goes back in a hard state, very similar to the one of the pre-outburst IBIS measurements (revolution 53), but in the lower energy range a disk black body component is still present. This behaviour is not unusual for BHC even if the physical reason is not fully understood (A. Zdziarski et al. 2004).

3.2. Spectral evolution of the source

The whole set of data has been fitted with standard XSPEC tools and the best fits have been obtained with a multicolor disk black body (Diskbb) model (Mitsuda et al. 1984) for the low energy data and a Thermal Comptonization model (CompTT) (Titarchuk et al. 1994) for the non thermal component. CompTT generally provides lower values for the plasma temperature kT_e if compared with other models (i.e. Compps) while modelling hard state spectra of BH transients, the advantage being the proper modelling of plasma with high optical thickness. The detailed spectral behaviour of the source is shown in Figure 4. At the beginning of the outburst (Rev53) the source is very weak and therefore the flux in the lower part of the JEM-X spectrum is below the instruments sensitivity. Therefore, it was possible to extract only a few points from JEM-X data in the range 15-22 keV and only a 2σ upper limit at lower energy. A single CompTT model fits the whole data set (15-150 keV) well, as shown in Table 1 with a $kT_e \sim 20$ keV and a $\tau_p \sim 3$. The point corresponding to revolution 53 in Figure 3 shows the correlation between the photon index and the flux for this data,

indicating a Low/Hard state (Grebenev et al. 2003). In a few days the source flux is then increasing quickly, and the statistics enable extraction of the SPI spectra as well (revolutions 56 and 60-61). The high energy spectral evolution clearly indicates that the source make a transition to the soft state as also shown in the HR evolution (see Figure 2 (c)). During revolution 56 the source was out of the JEM-X FOV therefore, only in this case, there are no data for $E < 20$ keV. But the source is quite luminous in the hard X-ray as shown by IBIS and, moreover, looking at the time spectral evolution (Figure 4), is possible to suppose that the source is still in a Low/Hard state. In revolutions 57-59 the 3-200 keV data can be fitted with a cutoff power law or with a thermal Comptonization Model without adding any black body component (Table 1). Conversely in revolutions 60-61 the disk black body appears very bright and the emission of this component account for 45% of the total luminosity. During the phase of decreasing flux (Rev 103), there was no evidence of a hard tail in the IBIS data (20-250 keV, 2σ upper limit corresponding to $0.2 \times 10^9 \text{erg} \times \text{cm}^{-2} \text{s}^{-1}$) while the source was clearly detected by JEM-X (see figure 4) and the data fitted with a Diskbb component showing a large value of the normalization constant (N_{Disk} , see Table 1), that is proportional to the square of the inner accretion disk radius (Mitsuda et al. 1984). The presence of a Diskbb in IGR J17464-3213 soft state is confirmed by the *RXTE* observation performed in 2003 May 28 (MJD=52787) (Homan et al. 2003). Two months later, close to the end of the outburst (revolutions 119-120), most of the flux is still due to the disk emission but a clear, though faint,comptonized high energy tail is present the IBIS data. The IBIS spectra taken at revolution 53 and 119-120,(i.e. at the beginning and at end of the outburst) are compatible with the same CompTT model parameters: it is clear that in both cases the source shows a hard tail in the high energy spectrum. Moreover, at the end of the outburst a disk component is still present, as also confirmed by the *RXTE* analysis of the data collected immediately before this *INTEGRAL* observation (from 2003 August 26 to 2003 September 23) (Lutovinov et al. 2004).

4. Discussion and Conclusion

The picture obtained from the spectral analysis in Figure 4 is quite clear, and summarised as follows. Phase 1: The source is in a “hard state”. At the beginning of the *INTEGRAL* monitoring campaign (revolution 53) the source has a low luminosity and the bulk of the emission is in the energy range 50-90 keV via Comptonisation of soft photons in an optically thick hot inner part of the disk or corona ($kT_e \sim 20$ keV and $\tau_p \sim 3$). In this phase there is no evidence of a soft disk component emission, with the disk being eventually truncated far from the hot inner region, though injecting soft photons that are up-scattered by the inner hot flow. In the next few days the amount of accreted matter is increasing

dramatically and in revolution 56 (~ 10 days later) the total emitted energy has increased by a factor of 30: the temperature of the Comptonised plasma is not changed and the optical thickness of the inner part of the corona become smaller, i.e. from $\tau_p \sim 3$ to $\tau_p \sim 1.3$ (revolution 57-59). There is still no evidence of soft emission from the system. Slowly, the peak emission shifts from 70 keV (revolution 53) to 40 keV (revolution 56) to finally 20 keV (revolution 57-59). The spectral evolution is shown in Figure 4 (top panel). Phase 2: The source is in a “soft/intermediate state” (revolution 60-61). Over a few days the average amount of energy released increases by a factor of 3 to 5, the optical thickness of the inner part of the corona becomes lower and lower ($\tau_p \sim 0.4$), the temperature of the hot Comptonised part increases accordingly up to ~ 40 keV and a bright soft disk component become visible, being responsible for the majority of the system energy release. The fit parameters are compatible with a T_{in} temperature of 1.5 keV and a disk close to the last stable orbit. The spectrum obtained with JEM-X, IBIS and SPI (Figure 4, middle panel) has a high statistical significance showing the presence of a substantial Comptonised component. A few days later the source has reached its maximum, showing a factor of two more flux but basically unchanged high energy behaviour (Parmar et al. 2003). Phase 3: The source enters the “soft state”. The last observations were performed a few months after the main peak of the outburst (revolution 103-120). The energy release has substantially decreased (factor of ~ 3), the hard emission has basically switched off, and the Comptonised photons energetics dropped by a factor of more than 10 with emission peaked at about 70 keV as in the pre-outburst state (revolution 53). This primarily indicates the temperature of the disk dropping to 0.9 keV, and the disk substantially outside from the last stable orbit. Soon afterwards, the source was no longer detectable by *INTEGRAL*; one month later it was also below the *ASM/RXTE* detection capability. The *INTEGRAL* observation of IGR J17464-3213 has been essential to disentangle the source behaviour and its spectral state evolution and, in turn, to better understand the physical processes active in the system in the different states.

We acknowledge the ASI financial/programmatic support via contracts I/R/389/02 and I/R/041/02; C. Spalletta for the careful editing of the manuscript, M. Federici for supervising the *INTEGRAL* data analysis system and A. Tarana for useful scientific discussions.

REFERENCES

- Courvoisier T.J.L., R. Walter, V. Beckmann, et al., A&A, 2003, 411,L53
Doxsey, H. Bradt, G. Fabbiano, et al., 1977, IAU Circ.3113
Goldwurm A., P. David, L. Foschini, et al., A&A, 2003, 411, L223

- Grebenev S. A., Lutovinov A. A., Sunyaev R. A. et al. 2003 Atel 189
- Homan J., Miller, J.M., Wijinands, R., et al. 2003 Atel 162
- Kaluzienski L.J. & Holt S.S., 1977, IAU Circ.3099
- Kretschmar, P., Chenevez, J., Capitanio, F., et al. 2003 Atel 180
- Lund N., C. Butz-Jorgensen, N.J. Westergaard, et al., A&A, 2003, 411,L231
- Lutovinov A., Revnivitsev M., S.Molkov et al. 2004 astro-ph/0407342
- Markwardt, C.B. & Swank,J.H. 2003, Atel 133
- Mitsuda K., Inoue H., Koyama K. et al. 1984, PASJ, 36, 741
- Parmar A.N., E. Kuulkers, T. Oosterbroek, et al., A&A, 2003, 411, L421
- Revnivitsev M., Chernyakova, M., Capitanio, F., et al. 2003, Atel 132
- Titarchuk L. 1994 ApJ,434,570
- Ubertini P., F. Lebrun, G. Di Cocco, et al., A&A, 2003, 411, L131
- Vedrenne, G., J.P. Roques, V. Shoenfelder, A&A 2003, 411, L63
- Winkler C. & T.L.L. Courvoisier, et al., A&A, 2003, 411, L1.
- A. Zdziarski & Gierlinski. Progress of theoretical Physics 2004 in press




Fig. 1.— *RXTE*-ASM one day averaged light curve. The gray dashed rectangles represent the simultaneous *INTEGRAL* observation periods corresponding to the spectra in figure 4. The time scale is represented in Modified Julian Date




Fig. 2.— (a) 20-30 keV and (b) 40-60 keV IBIS light curves from revolution 53 to revolution 61. The hardness ratio derived from the two energy ranges is shown in the bottom part of the figure (c)




Fig. 3.— Flux vs power law photon index for every SCW. The labels indicate the revolution number.




Fig. 4.— IBIS JEM-X and SPI common spectra in different states of the source outburst: the numbers on the top of each spectrum indicate the revolution number of the corresponding *INTEGRAL* observation.

Table 1. Fit parameter values (90 % confidence)

Revolution	T_{in} keV	N_{Disk}	kTe keV	τ_p	$N_{Comp} \times 10^2$	χ^2_ν	Deg.of freedom	flux ($\times 10^9$ $\text{erg} \times \text{cm}^{-2} \text{s}^{-1}$)	Fluxband keV
Rev53	...	...	20^{+15}_{-5}	3^{+4}_{-2}	$2.38^{+0.01}_{-0.01}$	0.7	7	~ 0.3	(15-250)
Rev56	...	...	24^{+3}_{-3}	$1.5^{+0.2}_{-0.3}$	$8.3^{+0.8}_{-0.7}$	0.96	14	~ 7	(20-250)
Rev57-59	...	...	19^{+5}_{-3}	$1.3^{+0.2}_{-0.4}$	$1.5^{+0.3}_{-0.3}$	1.03	57	~ 6	(3-250)
Rev60-61	$1.5^{+0.1}_{-0.1}$	77^{+16}_{-18}	33^{+14}_{-8}	$0.4^{+0.2}_{-0.2}$	8.3^{+2}_{-2}	1.01	106	~ 20	(3-250)
Rev 103	$1.09^{+0.02}_{-0.02}$	357^{+26}_{-39}	...	...	...	1.09	57	~ 3	(3-20)
Rev119-120	$0.87^{+0.02}_{-0.02}$	402^{+344}_{-214}	>13	<3	$1.5^{+5}_{-0.3}$	1.10	68	~ 1	(3-250)